

Barrandegnathus n. gen. (conodonta) from the Komstad Limestone (lower Mid Ordovician), Scandinavia, and its palaeogeographical significance

SVEND STOUGE



Stouge, S. 2005–12–31: *Barrandegnathus* n. gen. (conodonta) from the Komstad Limestone (lower Mid Ordovician), Scandinavia, and its palaeogeographical significance. *Bulletin of the Geological Society of Denmark*, Vol. 52, pp. 245–255. © 2005 by Geological Society of Denmark. ISSN 0011–6297.

A characteristic early Mid Ordovician conodont fauna first recorded from Bohemia is known from Morocco and southwestern Scandinavia. It is characterized by *Barrandegnathus* n. gen. (type species: *Baltoniodus bohemicus* Dzik, 1983). *Barrandegnathus bohemicus* is considered to be endemic to the Tornquist Sea, but it is often associated with Baltic Province conodonts.

The fauna was first described from the Klabava Formation the sediments of which accumulated in the Prague Basin, Czech Republic and was subsequently recorded as reworked Ordovician conodont fauna preserved in Upper Silurian strata from Morocco and from the Komstad Limestone in southwest Scandinavia. The stratigraphical range of the species is confined to the late part of the early Mid Ordovician (Darriwilian; Volkhov Regional Stage).

Barrandegnathus bohemicus probably migrated to Baltica with cold-water ocean currents moving from the high latitudinal Perunica terrane to the Gondwana supercontinent and to the southwestern margin of the Baltica continent.

Baltic Province conodonts are recorded from cool-water carbonates at or just off the periphery of Gondwana. Rifting along the Gondwana margin followed by drifting of palaeocontinents and terranes towards mid-high latitudes promoted deposition of cool-water carbonates. Baltic Province conodonts are recorded from Gondwana and Gondwana derived continents. The distribution of cool-water carbonates and the associated conodont fauna shows that Baltic Province conodonts were not confined to the Baltica continent.

Barrandegnathus n. gen. is introduced; the genus is represented by *B. bohemicus* (Dzik, 1983).

Keywords: Bornholm, Scania, conodonts, Middle Ordovician, biostratigraphy, palaeogeography, *Barrandegnathus* n. gen. *bohemicus* (Dzik, 1983).

Svend Stouge [svends@savik.geomus.ku.dk], Geological Museum, Øster Voldgade 5-7, DK-1350 Copenhagen K, Denmark.

Conodont faunas from relatively low yielding areas may be rewarding for correlative, provincial and palaeogeographic perspectives. One example of this is the conodont fauna first known from the Klabava Formation of Prague Basin, Czech Republic (Fig. 1; Dzik 1983a). The macrofauna of the Prague Basin is mostly endemic to the basin and hence fossil based correlation to other regions may be difficult (Havlíček & Fatka 1992). The Prague Basin is indeed not known for its abundance of conodonts, but the conodont biota reported from the succession i.e. *Baltoniodus bohemicus*, *Drepanodus arcuatus* and *Drepanoistodus basiovalis* (all *sensu* Dzik 1983a) is very characteristic. Dzik (1983a) placed this fauna within the

earliest Arenig and the conodonts were derived from the high palaeolatitudes on the southern hemisphere representing cold-water faunal association of the Mediterranean Province. The recent discovery of the same faunal elements associated with Baltic Province conodonts from the Khemis-n'Ga locality of western Morocco (Bultynck & Sarmiento 2003) indeed confirmed this idea as Morocco was situated on the Gondwana supercontinent and in a high latitudinal to nearly south polar location according to recent palaeogeographical models (Cocks & Torsvik 2002, 2004; Fortey & Cocks 2003).



Fig. 1. Map showing the locations with *Barrandegnathus* n. gen. (*) and Baltic Province conodonts (□) in Europe, Turkey and northwest Africa. The Baltic Province conodont fauna is found in Baltica, Gondwana (Morocco), (West) Avalonia (Ganderia-New Brunswick) and South China. *Barrandegnathus* is 'endemic' to the Tornquist Sea. The Baltic Province conodont fauna is found on Baltica and on continents that rifted and drifted away from Gondwana (i.e. Avalonia) and moved from high to mid high latitudes (or from low latitudes to mid latitudes i.e. South China) on the southern hemisphere.

The discovery of the same taxa from southwestern Scandinavia presented here does provide links from Perunica to Gondwana and Baltica. The record of the taxa in southwestern Scandinavia is from deep shelf environments. The fauna provides precise correlation between the deposits of the East European/Baltic Platform and the Prague Basin within the brief time interval of the early Mid Ordovician (i.e. within the upper Volkhov Regional Stage). In addition it provides new information to the previous ideas on faunal communication (e.g. Dzik 1983a, b; Gutiérrez-Marco *et al.* 1999) between the Perunica terrane, West Gondwana supercontinent and Baltica continent.

Material

The present material was isolated from the Komstad Limestone (Jaanusson 1960), which is known from

the Scania Region, Sweden and on Bornholm, Denmark (Figs 1, 2; Poulsen 1965, 1966; Bergström 1982; Stouge 1975; Nielsen 1995; Erlström *et al.* 2001; Stouge & Nielsen 2003) and in the German offshore G-14 well (Maletz 1997; Stouge 2001a).


The Komstad Limestone consists of lime mudstone and wackestone with thin silty partings and minor shale. The unit varies from a few meters to more than 15 m in thickness (Nielsen 1995, fig. 1).

The Komstad Limestone accumulated during the overall transgressive-regressive cycles of the Volkhov–Kundan stages (Nielsen 1992, 1995, 2004; Dronov & Holmer 1999; Laškovas 2000; Saadre *et al.* 2004).

Localities

The material described below was collected from the Limensgade and Skelbro localities on Bornholm,

Fig. 2. Location map with Komstad Limestone localities mentioned in the text. CDF = Caledonian Deformation Front; TESZ = Transeuropean Suture Zone.



Denmark, and from the Fågelsång locality and Albjåra well in Scania, Sweden (Fig. 2).

Skelbro and Limensgade, Bornholm

Location. Details on the location of the two Bornholm localities are given in Nielsen (1995; see Fig. 2).

Stratigraphy. Poulsen (1965, 1966) and Nielsen (1995) outlined the stratigraphy and described the trilobites from the Komstad Limestone on Bornholm. Stouge (1975) provided information on the conodont succession from the localities.

Lithological succession. The Komstad Limestone overlies the Mid Cambrian – Lower Ordovician Alum Shale Formation with a prominent disconformity. The

succession exposed consists from the bottom to the top of bedded limestone with minor glauconite and phosphate intraclasts - especially in the lower part - up to 1 m thick. Above follows a 2 m thick succession of two lithofacies. The lower 1 m consists of mudstone and separated by thin silt/shale. These are overlain by 1 m of wackestone with siltstone and marl.

About 1 m succession of wackestone to packstone concludes the Komstad Limestone on Bornholm. The uppermost part of the Komstad Limestone is phosphatised. On Bornholm, the *Dicellograptus* Shale formation disconformably overlies the Komstad Limestone.

Fauna and age of the succession. Trilobites from the Komstad Limestone are well documented. Brachiopods, ostracodes (Tinn & Meidla 1999) and cephalopods are also present. The trilobite assemblage is placed in the *Megistaspis polyphemus*, the *Megistaspis simon*, *Megistaspis limbata* trilobite zones of the Volk-

hov Regional Stage (Mid Ordovician) and the top of the limestone is referred to the *Asaphus expansus* trilobite Zone of the Kundan Stage (Nielsen 1995).

Conodonts are from the '*Baltoniodus*' *triangularis*, *Baltoniodus navis* and *Paroistodus originalis* zones (*sensu* Lindström 1971; Stouge 1975), the *Microzarkodina parva* (*sensu* Bagnoli & Stouge 1997), the *Baltoniodus norrlandicus* (*sensu* Stouge 1989) and the *Lenodus antivariabilis* (*sensu* Bagnoli & Stouge 1997) conodont zones of the Volkhov Regional Stage. The conodont assemblage from the top of the Komstad Limestone indicates the presence of the *Lenodus variabilis* Zone of the Kunda Stage.

Fågelsång, Scania, Sweden

Location. The precise location of the Fågelsång location (Fig. 2) is given by Ahlberg (1992), Nielsen (1995); Bergström *et al.* (2002) and Stouge & Nielsen (2003).

Stratigraphy. The Komstad Limestone overlies conformably the Tøyen Shale. About 3 m of the 5.3 m thick upper member are exposed (Stouge & Nielsen 2003). The upper part of the Komstad Limestone is exposed in old quarries at Fågelsång whereas the lower part is only known from wells (Nielsen 1995; Stouge & Nielsen 2003). The Almelund Shale conformably overlies the Komstad Limestone (Bergström *et al.* 2002).

Lithological succession. The exposed succession includes a wackestone unit with silty partings, a unit composed of nodular limestone and dark shale succeeded by a unit of wackestone and packstone. Bentonites are present in the succession. (Nielsen 1995; Stouge & Nielsen 2003).

Fauna and age of the Fågelsång succession. The lower member of the Komstad Limestone is referred to *Megistaspis simon* and probably includes lower *M. limbata* Subzone (Stouge & Nielsen 2003). The upper *Megistaspis limbata* Subzone is found in silty limestone changing upwards into cleaner wackestone below the nodular dark limestone and shale. The dark shale and nodular limestone belong to the *Asaphus expansus* Zone.

The conodont succession extends from *Baltoniodus norrlandicus* Zone and ranges into the lower part of the *Lenodus pseudoplanus* Zone (Stouge & Nielsen 2003, figs 5, 7).

Albjära well, western Scania, Sweden

Location. The Albjära well is situated in western Scania (see Lindholm 1991, fig 2; Fig. 2).

Stratigraphy. The Komstad Limestone in the Albjära-1 well is 3.5 m thick (Maletz 1995; Stouge *et al.* 2001). It is a distal exposure and much of the lower part of the Komstad Limestone is replaced by black shale referred to the Tøyen Shale. The Almelund Shale overlies the Komstad Limestone in the Albjära-1 well.

Lithological succession. The succession includes mudstone with minor shale partings in the lower part succeeded by an interval of nodular limestone in dark-grey shale. The Komstad Limestone and similar to Fågelsång concludes with a wackestone and packstone unit. Bentonites are preserved in the succession.

Fauna and age of the Albjära succession. Graptolites from the Tøyen Shale are from the *Undulograptus austrodentatus* (Arenig/Darriwillian) graptolite Zone and those from the overlying Almelund Shale are referred to the *Holmograptus lentus* Zone (Llanvirn; Maletz 1995, 2005; personal communication 2001). The conodonts are from the *Lenodus antivariabilis* Zone and extend into the *Lenodus pseudoplanus* Zone (Stouge *et al.* 2001).

Sedimentological and environmental interpretation of the Komstad Limestone

The sediments of the Komstad Limestone accumulated in deep shelf environments along the margin of the East European/Baltic platform (Stouge & Nielsen 2003).

The succession records the transition from carbonate accumulation of the deeper shelf to terrigenous sedimentation of the deep to very deep shelf with a return to carbonate accumulation of the shelf and to clastic deposition of the deep shelf.

Systematic palaeontology

All the specimens illustrated are housed in the collection of the Geological Museum, Copenhagen, and their catalogue numbers bearing the prefix MUGH.




Fig. 3. *Barrandegnathus bohemicus* (Dzik, 1983). All specimens are $\times 100$. – A. P1 element, MGUH 27756. Bornholm: Limensgade Quarry, # 55 – B. P1 element, MGUH 27757. Bornholm: Limensgade Quarry, # 56 – C. P2 element, MGUH 27758. Bornholm: Skelbro Quarry, # 9. – D. M element, MGUH 27759. Bornholm: Limensgade Quarry, # 55. – E. S0 element, MGUH 27760. Bornholm: Limensgade Quarry, # 55. – F. S1 element, MGUH 27761. Bornholm: Limensgade Quarry, # 56. – G. S2 element, MGUH 27762. Bornholm: Limensgade Quarry, # 55. – H. S3–4 element, MGUH 27763. Bornholm: Limensgade Quarry, # 56.

Family Balognathidae Hass, 1959

Genus *Barrandegnathus* n. gen.

Type species. *Baltoniodus bohemicus* Dzik, 1983a.

Derivation of name. After Joachim Barrande (1799–1883), the French palaeontologist that devoted his work to Bohemia.

Diagnosis. The septimembrate apparatus of *Barrandegnathus* is composed of P_1 , P_2 , M and a series of four morphological distinct S elements (*sensu* Purnell *et al.* 2000). The elements are predominantly hyaline with white matter present in the distal part of the denticles. Both P elements and the M element are robust with stout cusps and strong sharp edges. The P elements have short anterior and lateral, undenticulated to weakly denticulated processes.

Remarks. *Barrandegnathus* n. gen. resembles *Trapezognathus* Lindström, 1954, *Lenodus* Sergeeva, 1963 and *Eoplacognathus sensu* Dzik (1994) and Löfgren & Zhang (2003). *Barrandegnathus* is here considered related to the two first genera and hence it is placed in Balognathidae Hass, 1959 (*sensu* Stouge & Bagnoli 1999).

The *Barrandegnathus* P, M-element and the S_2 -element are diagnostic for the genus. The S_{3-4} -element of *Barrandegnathus* is not 'trapezoid' in shape, which otherwise is characteristic for the S_{3-4} -elements of *Trapezognathus* and *Lenodus*.

So far, only one species is referred to the new genus.

Barrandegnathus bohemicus (Dzik, 1983)
Fig. 3A–H

1983a *Baltoniodus bohemicus* sp. n., Dzik, pp. 339–342, pl. 2, fig 5; pl. 4, figs 1–4; pl. 5, figs 1–4.

2003 *Trapezognathus* sp., Bultynck & Sarmiento, pp. 267–268, pl. 2, figs. 5–6; pl. 3, figs. (?)1, 7, 9, 11, 13.

2003 *Trapezognathus quadrangulum* Lindström – Bultynck & Sarmiento, p. 267 (*partim*), pl. 4, fig 12.

Diagnosis. See diagnosis for the genus.

Remarks. Dzik (1983a) recorded and described all the elements of the species.

The M element resembles the corresponding element of *Gothodus*, Lindström in the development of a downwards directed and distally pointed anterior process. It differs by having a large stout cusp and it

System, series and stages		Baltoscandian Regional stages	Baltoscandian trilobite zones	Baltoscandian conodont zones		
ORDOVICIAN	MIDDLE	DARRIWILIAN	KUNDAN	<i>Asaphus raniceps</i>	<i>Lenodus pseudoplanus</i>	Range of <i>Barrandegnathus</i>
				<i>Asaphus expansus</i>	<i>Yangtzeplacognathus crassus</i>	
			VOLKHOV	<i>Megistaspis limbata</i>	<i>Lenodus antivariabilis</i>	
				<i>Megistaspis simon</i>	<i>Baltoniodus norrlandicus</i>	
				<i>Megistaspis polyphemus</i>	<i>Microzarkodina parva</i>	
					<i>Baltoniodus navis</i>	
					<i>Microzarkodina flabellum</i>	
					' <i>Baltoniodus</i> ' <i>triangularis</i>	

Fig. 4. Lower to Mid Ordovician conodont and trilobite zones of Baltoscandia and the range of *Barrandegnathus bohemicus* in the investigated sections. Conodont zones modified from Lindström (1971), Löfgren (1978, 1994, 1995); Bagnoli & Stouge (1997), Stouge & Bagnoli (1998) and Zhang (1998b).

may carry denticles on the upper margin of the base - a feature that otherwise characterizes the genera *Trapezognathus* and *Lenodus*.

The M element of *Trapezognathus* sp. illustrated by Bultynck & Sarmiento 2003 (pl. 3, fig. 1) probably is an M element of *Lenodus antivariabilis*. The denticulated element referred to *Trapezognathus quadrangulum* Lindström by Bultynck & Sarmiento (pl. 4, fig. 12) belongs to *Barrandegnathus* rather than to the *T. quadrangulum*.

Material. – 172 specimens (Skelbro: 57, Limensgade: 44, Fågelsång: 63; Albjära: 12).

Barrandegnathus n. gen. in the Komstad Limestone

Barrandegnathus bohemicus (Dzik, 1983) n. gen. characterises the fauna; it is associated with *Erraticodon* and '*Baltoniodus*' sp. (a prioniodontid genus composed of nondenticulated elements; see Stouge & Nielsen 2003, fig. 5). *Barrandegnathus* and '*Baltoniodus*' sp. are only known from the Komstad Limestone on Bornholm and in Scania. The two taxa are completely absent from the East European/Baltic Platform and have not been recorded from the Oslo Region (Rasmussen 1991, 2001).

Other common associated species include *Baltoniodus norrlandicus*, *Lenodus antivariabilis*, *Microzarkodina parva*, *Protopanderodus* spp., *Scalpellodus* and *Cor-*

nuodus. Species of *Drepanoistodus* and *Drepanodus* are also present. These taxa are together with *Erraticodon* more widespread and occur on the platform and platform margin of Baltica (e.g. Bednarczyk 1998; Löfgren 1978, 1995, 2000a, b; Rasmussen 1991, 2001; Viira *et al.* 2001; Zhang 1998b) and on other palaeocontinents (Nowlan 1981; Landing *et al.* 2003). The associated taxa are (except for *Erraticodon*) considered typical of the Baltic Province of the Atlantic Realm (Bagnoli & Stouge 1997; Rasmussen 1998).

Barrandegnathus bohemicus (Dzik 1983) n. gen. and the most characteristic associated species in the Komstad Limestone appear within the silty limestone lithofacies and extend into the succeeding wackestone. The taxon does not extend into the uppermost part of the Komstad Limestone (Fig. 4; Stouge & Nielsen 2003, fig. 5).

Comparison of *Barrandegnathus bohemicus*-bearing strata

Bornholm and Scania. On Bornholm, Denmark and Scania, Sweden the range of *Barrandegnathus bohemicus* is from the top of *Baltoniodus norrlandicus* Zone to near the top of the *Lenodus antivariabilis* Zone (Fig. 4). The species is constrained to the *Megistaspis limbata* Zone of the upper Volhov Regional Stage (Fig. 4; Nielsen 1995; Stouge & Nielsen 2003).

In Scandinavia, the *Megistaspis limbata* Zone corresponds to the lowermost part of the Darriwilian Stage

of the international Middle Ordovician Series (Stouge & Nielsen 2003).

Czech Republic. The sediments of the Prague Basin are very rich in fossils, and the macrofauna and microfauna/flora are referred to the Mediterranean Province of the high latitude on the southern hemisphere (Havlíček 1989; Havlíček *et al.* 1994).

The low-diversity conodont fauna (Dzik 1983a) was recovered from siliciclastic sediments associated with Fe-nodules and volcanics, which are referred to the Klabava Formation (Havlíček 1981; Havlíček & Fatka 1992). *Barrandegnathus bohemicus* occurs in the *Azygograptus ellesi-Tetragraptus reclinatus abbreviatus* graptolite Zone (*sensu* Kraft & Kraft 1999), which correlates with the upper part of the Arenig (Bohemia/Gondwana) Series.

Morocco. The *Barrandegnathus* fauna from the Khemiss'n'Ga locality in western Morocco was recorded as reworked Ordovician conodont fauna from an Upper Silurian silty-carbonate unit (Bultynck & Sarmiento 2003). The Silurian sediments overlie lower – mid Arenigian sand and siltstones with a prominent erosional unconformity (Beun *et al.* 1986). The conodont fauna is the first record of early Mid Ordovician faunas from western Morocco, thus representing an unexposed interval in the region (Bultynck & Sarmiento 2003; see also Gutiérrez-Marco *et al.* 2003). The Moroccan conodont fauna is relatively diverse and resembles in most aspects the Komstad Limestone fauna of southwestern margin of Baltica – in the presence of *Barrandegnathus* but also with the additional presence of other Baltic taxa – allowing for a correlation to the *Baltoniodus norrlandicus* and *Lenodus antivariabilis* conodont zones.

Palaeogeographical distribution of *Barrandegnathus*

Barrandegnathus is recorded from the Prague Basin, which belongs to the single peripheral terrane Perunica of the West Gondwana supercontinent, West Gondwana supercontinent and the southwestern margin of Baltica continent (Havlíček 1981, 1989; Havlíček *et al.* 1994; Cocks & Torsvik 2002, 2004; Fortey & Cocks 2003). All three regions were facing the Tornquist Sea (Cocks & Torsvik 2002, 2004; Fortey & Cocks 2003; Torsvik & Rehnström 2003). The Perunica terrane maintained a high latitudinal position, whereas Morocco (West Gondwana) with its occurrence of *Barrandegnathus* and the Baltic fauna was

situated in a high to mid-high latitudinal position during the lower Mid Ordovician. *Barrandegnathus* was restricted to the Tornquist Sea and Perunica and Morocco were close to each other during late Arenig. The Komstad Limestone accumulated at a mid-southerly position on the Baltica palaeocontinent and was separated from Perunica and Gondwana by the Tornquist Sea (Cocks & Torsvik 2002, 2004; Fortey & Cocks 2003; Torsvik & Rehnström 2003). On Baltica *Barrandegnathus* is – as mentioned above – only found in the Komstad Limestone.

Distribution of Baltoscandian conodonts adjacent to Gondwana

Other terranes peripheral to West Gondwana i.e. Armorica and Iberia (Fortey & Cocks 2003; Gutiérrez-Marco *et al.* 1999, 2003) have not yielded conodonts contemporaneous to the *Barrandegnathus* fauna. The areas Turkey, South China and New Brunswick, Canada however provide further information.

Turkey. Early Darriwilian conodonts are known from several localities in the Taurides or the Tauride Belt (Barnes *in* Dean 1973; Gedik 1977; Sarmiento *et al.* 1999). The Cambrian – Mid Ordovician Seydisehir Formation is predominantly a siliciclastic unit with few nodular limestone beds. The sediments accumulated on the extensive Tauride Platform at the Gondwana margin but two distinct limestone units (i.e. the Sobova Limestone and the younger Tekmen Member) interrupt the overall siliciclastic sedimentation of the Seydisehir Formation (Gedik 1977; Sarmiento *et al.* 1999; Kozlu *et al.* 2003).

The yield of conodonts is relatively low in the sand and siltstones but the limestone nodules and the two limestone units produced early and mid Ordovician conodonts that in general show Baltic Province affinities. One exception to this is the report of *Oepikodus evae* (= *Prioniodus evae* of Gedik 1977) and *Bergstroemognathus* sp. (i.e. *Falodus* sp. of Gedik 1977) – a faunal association, which is more typical for low latitudinal East Gondwana and Laurentia rather than Baltica.

The well-developed Mid Ordovician conodont fauna recorded from the Sobova Limestone (Barnes *in* Dean 1973; Gedik 1977) shows close affinities with the fauna from the East European/Baltic Platform. The fauna from the Sobova Limestone correlates to the Baltoscandian *Baltoniodus norrlandicus* and *Lenodus antivariabilis* conodont zones.

South China. The Dawan Formation in the Hubei district of South China consists of a complex mixture of carbonate, shale and cool-water carbonates. The associated conodont fauna changes from warm-water types to conodonts with Baltic affinity (An 1981, 1987; An *et al.* 1985; Lindström *et al.* 1991; Stouge 2001b; Zhang 1998a). The faunal shift occurs in the late Early Ordovician and the base of the Middle Ordovician is characterized by the appearance of '*Baltoniodus*' *triangularis* (Wang *et al.* 2005). From this level the succession lithologically resembles that of Baltoscandia (Lindström *et al.* 1991) and the conodont zones can be matched to those known from Baltoscandia (Stouge 2003). The *Baltoniodus norrlandicus*, *Lenodus antiovariabilis* and younger zones can be distinguished and the fauna is clearly of Baltoscandian affinity (Zhang 1998a; Wang & Bergström 1999; Stouge 2003).

New Brunswick. Mid – late Arenigian conodonts with strong Baltic Province affinity are known from two locations in New Brunswick, Canada (Nowlan 1981; Landing *et al.* 2003) i.e. the (west) Avalonia (= Ganderia microcontinent of van Staal *et al.* 2003). The first occurrence is from central New Brunswick, where *Microzarkodina flabellum* s.l. is recorded from the uppermost part of the lower Tetagouche Group.

A Mid Ordovician (*Lenodus antiovariabilis* and *L. variabilis* zones) conodont fauna is recorded from a limestone boulder in the Triassic Lepreau Formation in southern New Brunswick. The conodont yielding limestone unit is a cool-water carbonate with a diverse conodont fauna of Baltic affinity (Landing *et al.* 2003). The appearance of the conodonts of Baltic Province suggests that west Avalonia (i.e. Ganderia) reached a high to mid latitudinal position (e.g. Landing 1998, 2003) during the late Arenig.

Neither of these pre-Caradoc Baltic conodont faunas is associated with *Barrandegnathus*.

Discussion

The arrival of *Barrandegnathus* in late Volkhov to Baltica was probably a result of various palaeobiogeographical factors, notably the extent and timing of the late Volkhov transgressive-regressive event within the Tornquist Sea area. Tectonic movements near West Gondwana and Perunica possibly associated with rifting opened for the northern direction of a cold high latitudinal ocean current (see Christiansen & Stouge 1999a, b). This event promoted the emigration of the Perunica/Gondwana-derived conodonts to Baltica in the late Volkhov times.

The ocean current thus provided a bridge for the conodont species to migrate across the Tornquist Sea and the cold-water 'endemic' *Barrandegnathus bohemicus* was not exclusively constrained to high latitudinal sites.

The cold oceanic current remained active long enough for *B. bohemicus* to colonize the cool and deep-water environments along the southwestern margin of Baltica. Towards the end of Volkhov times this route was closed and the Gondwana and Baltica populations became isolated from each other again.

Available evidence shows that Baltoscandian conodonts (or the Baltic Province) were not constrained to the Baltica continent during early Mid Ordovician times. The Baltoscandian conodont fauna was associated with cool-temperate water carbonate depositional environments (e.g. Lindström 1984a, b). Apparently, cool-water carbonates accumulated on Gondwana i.e. Morocco and on Gondwana derived microcontinents (South China, New Brunswick) as they reached mid-high latitudes. The associated Baltic Province conodont fauna inhabited the continents as these reached or passed through distinct latitudinal and climate belts on the southern hemisphere.

The early Darriwilian conodont faunas from Turkey, New Brunswick and South China are all similar and this suggests that the areas occupied a relatively lower northern position than Perunica did during the lower Mid Ordovician. Apparently major changes in the oceanic current system occurred at this time allowing for cool-water carbonate accumulation at places such as Morocco and Turkey. According to recent palaeogeographical models these areas were situated near the south pole (Cocks & Torsvik 2002, 2004; Fortey & Cocks 2003).

Conclusions

Barrandegnathus originated from west Gondwana and occupied a restricted area within the Tornquist Sea between western Morocco on Gondwana and Perunica, and at high latitudes on the southern hemisphere. *Barrandegnathus* is recorded from the Komstad Limestone and the taxon lived in deep-water shelf environments along the margin of Baltica. It reached Baltica and stayed on its continental margin within a relative short time interval of the early Mid Ordovician (early Darriwilian, late Volkhov).

The immigration of *Barrandegnathus* to southwestern Baltica was probably due to the presence of cold-water oceanic currents (e.g. Dzik 1983a; Christiansen & Stouge 1999a, b). Its arrival to Baltica marks the base of the Darriwilian international Stage of the glo-

bal Middle Ordovician Series (Mitchell & Maletz 1995; Stouge & Nielsen 2003).

Barrandegnathus bohemicus has not been recorded from other Gondwana-derived terranes. However, conodonts with Baltoscandian affinities and contemporaneous to the *Barrandegnathus* fauna appeared near to or along the margin of Gondwana suggesting that faunal communication occurred between Gondwana, peri-Gondwana and Baltica. The Baltic Province conodont fauna also appeared on continents (i.e. Avalonia) that drifted northwards (New Brunswick-Turkey) and southwards (South China). The appearance of Baltic Province conodonts at these places was probably a consequence of the drifting of the palaeocontinents towards mid latitudinal sites with distinct temperate climate belts.

Acknowledgements

J. Halskov drafted the figures. H.J. Hansen helped with the SEM pictures. J. A. Rasmussen read an early version and NN corrected the English language. The Bulletin referees M. Lindström and G. Sarmiento are gratefully thanked for many helpful comments. The Danish Natural Science Research Council financially supported the work on Bornholm and in Scania.

References

- Ahlberg, P. 1992: Fågelsångsdalen – et klassisk exkursionsområde. Lunds Geologiska Fältklubbs jubileumsskrift 1892–1992, 27–37.
- An, Tai-Xiang 1981: Recent progress in Cambrian and Ordovician conodont biostratigraphy of China. Geological Society of America Special Paper 187, 209–226.
- An, Tai-Xiang 1987: The Lower Paleozoic conodonts of South China. Beijing University Press, Beijing (in Chinese), 238 pp. (in Chinese).
- An, Taixiang, Du Gouqing & Gao Qinqin 1985: Ordovician conodonts from Hubei, China. Geological publishing House, Beijing. 64 pp. (in Chinese).
- Bagnoli, G. & Stouge, S. 1997: Lower Ordovician (Billingenian–Kundan) conodont zonation and provinces based on sections from Horns Udde, north Öland, Sweden. Bolletino della Società Paleontologica Italiana 20, 109–163.
- Bednarczyk, W.S. 1998: Ordovician conodont biostratigraphy of the Polish part of the Baltic Syncline. In: Szaniawski, H. (ed.): Proceedings of the Sixth Conodont Symposium (ECOS VI). Palaeontologia Polonica 58, 107–121.
- Bergström, J. 1982: Scania. Field Excursion Guide IV Intern. In: Bruton, D.L. & Williams, S.H. (eds): Symposium on the Ordovician System. Palaeontological Contributions for the University of Oslo 279, 184–191.
- Bergström, S.M., Larsson, K., Pålsson, C. & Ahlberg, P. 2002: The Almelund Shale, a replacement name for the Upper *Didymograptus* Shale and the Lower *Dicellograptus* Shale in the lithostratigraphical classification of the Ordovician succession in Scania, Southern Sweden. Bulletin of the Geological Society of Denmark 49, 41–48.
- Beun, N., Boulanger, D., Huvelin, P. & Tajeddine, K. 1986: Le Paléozoïque de Khemis-n’Ga: un Paléorelief sous la série gypseuse jurassica–crétacée des Mouissat (région de Safi, Maroc). Comptes Rendus de l’Académie des Sciences, Paris 302, 39–42.
- Bultynck, P. & Sarmiento, G. N. 2003: Reworked Ordovician and autochthonous Siluro–Devonian conodonts from Khemis-n’Ga (Moroccan Meseta) – Depositional, environmental, and palaeogeographic implications. Courier Forschungs Institut Senckenberg 242, 257–283.
- Christiansen, J. & Stouge, S. 1999a: Oceanic circulation as an element in palaeogeographical reconstructions: the Arenig (early Ordovician) as an example. Terra Nova 11, 73–78.
- Christiansen, J. & Stouge, S. 1999b: Using palaeo-oceanographical modelling in reconstruction Early Ordovician palaeogeography. In: Kraft, P. & Fatka, O. (eds): *Quo vadis Ordovician?* Acta Universitatis Carolinae, Geologica 43, 515–518.
- Cocks, L.R.M. & Torsvik, T.H. 2002: Earth geography from 500 to 400 million years ago: a faunal and palaeomagnetic review. Journal of the Geological Society, London 159, 631–644.
- Cocks, L.R.M. & Torsvik, T.H. 2004: Major terranes in the Ordovician, 61–67. In: Webby, D.D., Paris, F., Droser, M.L. & Percival I.G. (eds): The great Ordovician biodiversification event. Columbia University Press, New York.
- Dean, W.T. 1973: The Lower Palaeozoic stratigraphy and faunas of the Taurus Mountains near Beysehir, Turkey. III. The trilobites of the Sobova Formation. Bulletin British Museum (Natural History), Geology 24, 279–348.
- Dronov, A. & Holmer, L. 1999: Depositional sequences in the Ordovician of Baltoscandia. In: Kraft, P. & Fatka, O. (eds): *Quo vadis Ordovician?* Acta Universitatis Carolinae, Geologica 43, 133–138.
- Dzik, J. 1983a: Early Ordovician conodonts from the Barrandian and Bohemian-Baltic faunal relationships. Acta Palaeontologica Polonica 28 (3–4), 327–368.
- Dzik, J. 1983b: Relationships between Ordovician Baltic and North American Midcontinent conodont faunas. Fossils and Strata 15, 59–85.
- Dzik, J. 1994: Conodonts from the Mójca Limestone. Palaeontologia Polonica 53, 43–128.
- Earlström, M., Ahlberg, P. & Löfgren, A. 2001: Lower Palaeozoic stratigraphy at Lyby and Tängelsås, central Scania, southern Sweden. GFF 123, 7–14.
- Fortey, R.A. & Cocks, R.M. 2003. Palaeontological evidence bearing on global Ordovician–Silurian continental reconstructions. Earth-Science Reviews 61, 245–307.
- Gedik, I. 1977: Conodont biostratigraphy in the middle Taurus. Geological Society of Turkey Bulletin 20, 35–48.
- Guetiérrez-Marco, J.C., Rábano, I., Sarmiento, G.N., Aceñola, G.F., José, M.A. s., Pieren, A.P., Herranz, P., Couto, H.M., Piçarra, J.M. 1999: Faunal dynamics between Iberia and Bohemia during the Oretanian and Dobrotivian (late Middle – earliest Upper Ordovician) and biogeographic relations with Avalonia and Baltica. In: Kraft, P. & Fatka, O. (eds): *Quo vadis Ordovician?* Acta Universitatis Carolinae, Geologica 43, 487–490.

- Gutiérrez-Marco, J.C., Destombes, J., Rábano, I., Aceñolaza, G.F., Sarmiento, G.N. & José, M.Á.S. 2003: El Ordovícico Medio del Anti-Atlas marroquí: paleobiodiversidad, actualización bioestratigráfica y correlación. *Geobios* 36, 151–177.
- Hass, W.H. 1959: Conodonts from the Shappel Limestone of Texas. U.S. Geological Survey Professional Paper 294J, 365–400.
- Havlíček, V. 1981: Development of a linear sedimentary depression exemplified by the Prague Basin (Ordovician – Middle Devonian, Barrandian area – central Bohemia). *Sborník Geologických ved. Geologie* 35, 7–48.
- Havlíček, V. 1989: Climatic changes and development of benthic communities through the Mediterranean Ordovician. *Sborník geologických ved. Geologie* 44, 79–116.
- Havlíček, V. & Fatka, O. 1992: Ordovician of the Prague Basin (Barrandian area, Czechoslovakia). In: Webby, D.D. & Laurie, J.R. (eds): *Global perspectives on Ordovician geology*, 461–472. Proceedings of the sixth international symposium on the Ordovician System, University of Sydney, Australia. A.A. Balkema, Rotterdam.
- Havlíček, V., Vanek, J. & Fatka, O. 1994: Perunica microcontinent in the Ordovician (its position with the Mediterranean Province, series division, benthic and pelagic associations). *Sborník geologických ved. Geologie* 46, 25–56.
- Jaanusson, V. 1960: Graptoloids from the Ontikan and Viruan (Ordov.) limestones of Estonia and Sweden. *Bulletin of the Geological Institutions of the University of Upsala* 38, 289–366.
- Kozlu, H., Göncüoğlu, Sarmiento, G. N. & Gül, M.A. 2002: Mid-Ordovician (Late Darriwilian) conodonts from the southern-central Taurides: geological implications. *Turkish Journal of Earth Sciences* 11, 113–126.
- Kraft, J. & Kraft, P. 1999: Graptolite biostratigraphy of the Lower and Middle Ordovician of Bohemia. In: Kraft, P. & Fatka, O. (eds): *Quo vadis Ordovician?* Acta Universitatis Carolinae-Geologica 43, 33–36.
- Landing, E. 1998: Avalon and Gondwana – the antipodean dance of two Early Paleozoic continents. *Geological Society of America, Abstracts with Programs* 30 (7), 18.
- Landing, E. 2003: Avalon – non-Gondwanan microcontinent by the late Precambrian: new evidence from the Precambrian–Cambrian boundary interval. *Geological Society of America, Abstracts with Programs* 35 (3), 18.
- Landing, E., Westrop S. R. & Dong Hee Kim 2003: First Middle Ordovician biota from southern New Brunswick: stratigraphic and tectonic implications for the evolution of the Avalon continent. *Canadian Journal of Earth Sciences* 40, 715–730.
- Laškovas, J. 2000: The sedimentation environments of the Ordovician basin in the south-western margin of the East European Platform and lithogenesis of deposits. 314 pp. Institute of Geology, Vilnius.
- Lindholm, K. 1991: Hunnebergian graptolites and biostratigraphy in southern Scandinavia. *Lund Publications on Geology* 95, 36 pp.
- Lindström, M. 1954: Conodonts from the lowermost Ordovician strata of South-Central Sweden. *Geologiska Föreningens i Stockholm Förhandlingar* 76, 517–604.
- Lindström, M. 1971: Lower Ordovician conodonts of Europe. *Geological Society of America Memoir* 127, 21–61.
- Lindström, M. 1984a: The Ordovician climate based on the study of carbonate rocks, 81–88. In: Bruton D.L. (ed.): *Aspects of the Ordovician System*. Palaeontological Contribution from the University of Oslo 295, 81–88. Universitetsforlaget, Oslo.
- Lindström, M. 1984b: Baltoscandic conodont life environments in the Ordovician: Sedimentologic and paleogeographic evidence. *Geological Society of America, Special Paper* 196, 33–42.
- Lindström, M., Chen Jun-Yuan & Zhang Jun-Min 1991: Section at Daping reveals a Sino-Baltoscandian parallelism of facies in the Ordovician. *Geologiska Föreningens i Stockholm Förhandlingar* 113, 189–205.
- Löfgren, A. 1978: Arenigian and Llanvirnian conodonts from Jämtland, northern Sweden. *Fossils and Strata* 13, 129 pp.
- Löfgren, A. 1994: Arenig (Lower Ordovician) conodonts and biozonation in the eastern Siljan District, central Sweden. *Journal of Paleontology* 68, 1350–1468.
- Löfgren, A. 1995: The middle Lanna/Volkhov Stage (middle Arenig) of Sweden and its conodont fauna. *Geological Magazine* 132, 693–711.
- Löfgren, A. 2000a: Conodont biozonation in the upper Arenig of Sweden. *Geological Magazine* 137, 53–65.
- Löfgren, A. 2000b: Early to early Middle Ordovician conodont biostratigraphy of the Gillberga quarry, northern Öland, Sweden. *GFF* 122, 321–338.
- Löfgren, A. & Zhang, J. 2003: Element association and morphology in some Middle Ordovician platform-equipped conodonts. *Journal of Paleontology* 77, 723–739.
- Maletz, J. 1995: The Middle Ordovician (Llanvirn) graptolite succession of the Albjärna core (Scania, Sweden) and its implication for a revised biozonation. *Zeitschrift für geologische Wissenschaften* 23, 249–259.
- Maletz, J. 1997: Ordovician and Silurian strata of the G-14 well (Baltic Sea): graptolite faunas and biostratigraphy. *Zeitschrift für geologische Wissenschaften* 25, 29–39.
- Maletz, J. 2005: Early Middle Ordovician graptolite biostratigraphy of the Lovisefred and Albjärna drill cores (Scania, southern Sweden). *Palaeontology* 48 (3), 763–780.
- Mitchell, C.E. & Maletz, J. 1995: Proposal for adoption of the base of the *Undulograptus austrodentatus* Biozone as a global Ordovician stage and series boundary level. *Lethaia* 28, 317–331.
- Nielsen, A.T. 1992: Ecostratigraphy and the recognition of Arenigian (Early Ordovician) sea-level changes, 355–366. In: Webby, B.D. & Laurie, J.R. (eds): *Global Perspectives on Ordovician Geology*. Balkema, Rotterdam.
- Nielsen, A.T. 1995: Trilobite biostratigraphy, palaeoecology and systematics of the Komstad Limestone and Huk Formation (Lower Ordovician), southern Scandinavian. *Fossils and Strata* 38, 374 pp.
- Nielsen, A.T. 2004: Ordovician sea level changes: A Baltoscandian perspective, 84–93. In: Webby, B.D., Paris, F., Droser, M.L. & Percival, I.G.: *The great Ordovician biodiversification event*. Columbia University Press, New York.
- Nowlan, G. 1981: Some Ordovician conodont faunules from the Miramichai Anticlinorium, New Brunswick. *Geological Survey of Canada, Bulletin* 345, 35 pp.
- Poulsen, V. 1965: An Early Ordovician trilobite fauna from Bornholm. *Meddelelser fra Dansk Geologisk Forening* 16, 49–113.
- Poulsen, V. 1966: Cambro–Silurian stratigraphy of Bornholm. *Meddelelser fra Dansk Geologisk Forening* 16, 117–137.

- Purnell, M.A., Donoghue, P.C.J. & Aldridge, E. 2000: Orientation and anatomical notation in conodonts. *Journal of Paleontology* 74, 113–122.
- Rasmussen, J.A. 1991: Conodont stratigraphy of the lower Ordovician Huk Formation at Slemmestad, southern Norway. *Norsk Geologisk Tidsskrift* 71, 265–288.
- Rasmussen, J.A. 1998: A reinterpretation of the conodont Atlantic Realm in the late Early Ordovician (early Llanvirn). In: Szaniawski, H. (ed.): *Proceedings of the Sixth European Conodont Symposium (ECOS VI)*. *Palaeontologica Polonica*, 67–77.
- Rasmussen, J.A. 2001: Conodont biostratigraphy and taxonomy of the Ordovician shelf margin deposits in the Scandinavian Caledonides. *Fossils and Strata* 48, 180 pp.
- Saadre, T., Einasto, R., Nolvak, J. & Stouge, S. 2004: Ordovician stratigraphy of the Kovel-1 well (Volkhov–Haljala) in the Volynia region, northwestern Ukraine, *Bulletin of the Geological Society of Denmark* 51, 47–69.
- Sarmiento, G.N., Gül, M.A., Kozlu, H & Göncüoğlu, M.C. 1999: Darriwilian conodonts from the Taurus Mountains, Southern Turkey. In: Kraft, P. & Fatka, O. (eds): *Quo vadis Ordovician?* *Acta Universitatis Carolinae-Geologica* 43, 37–40.
- Sergeeva, S.P. 1963: A new Early Ordovician conodont genus of the family Prioniodinidae. *Academy Nauk. SSR, Palaeontological Journal* 1963, 138–140.
- Stouge, S. 1975: Conodontzonerne i orthoceratitkalken (Nedre Ordovicium) på Bornholm og i Fågelsång. *Dansk Geologisk Forening, Årsskrift* 1974, 32–38.
- Stouge, S. 1989: Lower Ordovician (Ontikan) conodont biostratigraphy in Scandinavia. *Proceedings of the Academy of Sciences of the Estonia SSR Geologia* 38, 68–72.
- Stouge, S. 2001a: Lower Ordovician conodonts from the G-14 well, Baltic Sea (Germany). In: Giese, U., Katzung, G. & Miller (eds): *The Rügen Caledonides-southern Baltic Sea*. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 222, 141–160.
- Stouge, S. 2001b: Stratigraphic and biogeographic affinities of the Lower – Middle Ordovician conodonts of the Yangtze Platform, South China, 27. In: Harper, D.A.T. (ed.): *45th Annual Meeting of the Palaeontological Association*. Abstracts. Geological Museum. University of Copenhagen. Copenhagen.
- Stouge, S. & Bagnoli, G. 1989: Lower Ordovician (Volkhovian–Kundan) conodonts from Hagudden, northern Öland, Sweden. *Paleontographica Italica* 77, 1–54.
- Stouge, S. & Bagnoli 1999: The suprageneric classification of some Ordovician prioniodontid conodonts. *Bollettino della Società Paleontologica Italiana* 37, 145–158.
- Stouge, S. & Nielsen, A.T. 2003: An integrated biostratigraphical analysis of the Volkhov–Kunda (Lower Ordovician) succession at Fågelsång, Scania, Sweden. *Bulletin of the Geological Society of Denmark* 50, 75–94.
- Stouge, S., Maletz, J. & Nielsen, A.T. 2001: Integrated biostratigraphy (graptolites, trilobites and conodonts) and interregional correlation of the upper Volkhov – lower Kunda deposits of Albjära-1 well, Scania, Sweden, 27–28. In: Harper, D.A.T. & Stouge, S. (eds): *WOGOGO-2001*. Working Group on the Ordovician Geology of Baltoscandia. Øresund Region, mid-May 2001. Abstracts. Geological Museum, University of Copenhagen, Copenhagen.
- Tinn, O. & Meidla, T. 1999: Ordovician ostracodes from the Komstad Limestone. *Bulletin of the Geological Society of Denmark* 46, 25–30.
- Torsvik, T.H. & Rehnström, E.F. 2003: The Tornquist Sea and Baltica-Avalonia docking. *Tectonophysics* 362, 67–82.
- van Staal, C.R., Wilson, R.A., Rogers, N., Fyffe, L.R., Langton, J.P., McCutcheon, S., McNicoll, V. & Ravenhurst, C.E. 2003: Geology and tectonic history of the Bathurst Supergroup, Bathurst Mining Camp, and its relationships to coeval rocks in southwestern New Brunswick and adjacent Maine – a synthesis. *Economic Geology, Monograph* 11, 37–60.
- Viira, V., Löfgren, A., Mägi, S. & Wickström, J. 2001: An Early to Middle Ordovician succession of conodont faunas at Mäekalda, northern Estonia. *Geological Magazine* 138, 699–718.
- Wang, Z.H. & Bergström, S.M. 1999: Conodonts across the base of the Darriwilian Stage in South China. *Acta Micropalaeontologica Sinica* 16, 325–350.
- Wang, X., Stouge, S., Erdtmann, B.-D., Chen, Z., Li, C., Wang, Q., Zeng, Z. & Chen, H. 2005: A proposed GSSP for the base of the Middle Ordovician Series: the Huanghuachang section, Yichang, China. *Episodes* 28, 105–118.
- Zhang, J. 1998a: Conodonts from the Guniutan Formation (Llanvirnian) in Hubei and Hunan Provinces, south-central China. *Stockholm Contributions in Geology* 46, 161 pp.
- Zhang, J. 1998b: The Middle Ordovician conodonts from the Atlantic Faunal Region and the evolution of key conodont genera. *Meddelanden från Stockholms Universitets Institution för Geologi och Geokemi* 298, 5–27.

